

- State Mus. Ann. Rept.*, 28: 48. (Title-page date, 1879, seems to be an error.)
- Pennington, L. H. 1915. *Marasmius*. *N. A. Flora*, 9: 250-286.
- Seaver, F. J. 1928. *The North American Cup-Fungi*. Pp. 3-284.
- Smith, Alexander H. 1935a. Unusual Agarics from Michigan, II. *Mich. Acad. Sci., Arts and Letters*, 20: 171-183.
- Smith, Alexander H. 1935b. Studies in the Genus *Mycena*, II. *Mycologia*, 27: 586-604.
- Smith, Alexander H., and L. E. Wehmeyer. 1936. Contributions to a Study of the Fungous Flora of Nova Scotia, II. Agaricaceae and Boletaceae. *Mich. Acad. Sci., Arts and Letters*, 21: 163-197.
- Snell, W. H. 1936. Notes on *Boletus*, V. *Mycologia*, 28: 463-475.

VETERINARIANS O. K. REMEDY FOR HEART WORMS IN DOGS

The comparatively new treatment for heart-worm infestation in dogs has given good results in most cases when used by practicing veterinarians according to a check made recently by zoologists of the United States Bureau of Animal Industry.

The treatment, which was perfected by zoologists of the Bureau and has been used for more than a year, consists of injections into the blood stream of the drug known as Fuadin. Dogs affected with heart worms may become greatly exhausted when hunting and many of them die.

In checking the effectiveness of the remedy, the Bureau's study showed that out of a total of 1,000 dogs treated by the method, more than 90 per cent were restored to usefulness. Blood tests of dogs that had received treatment showed that the microscopic parasite had disappeared from the blood stream. The remedy is therefore considered the most practical means of treating heart-worm disease in dogs.