

- Hubbs, Carl L., and Laura C. 1933. Increased Growth, Predominant Maleness and Apparent Infertility of Hybrid Sunfishes. *Pap. Mich. Acad. Sci., Arts and Letters*, 17 (1932): 613-641.
- Hubbs, Carl L., and Gerald P. Cooper. 1936. Minnows of Michigan. *Cranbrook Institute of Science*, Bull. 8: 195.
- Langlois, T. H. 1936. A Study of the Small-Mouth Bass, *Micropterus Dolomieu* (Lacepede), in Rearing Ponds in Ohio. *Ohio Biol. Surv.*, Bull. 33: 191-225.
- Potter, George E. 1925. Scales of the Bluegill, *Lepomis pallidus* (Mitchell). *Trans. Am. Microsc. Soc.*, 44: 31-37.
- Tester, Albert L. 1932. Rate of Growth of the Small-Mouthed Black Bass (*Micropterus dolomieu*) in Some Ontario Waters. *Publ. Ont. Fish. Res. Lab.*, 47: 205-221.
- Thompson, David H., and Donald F. Hansen. 1937. Research on the Large-Mouth Black Bass. *Ill. Conservation*, 2 (No. 2, Spring): 1-24.
- Wright, Stillman. 1929. A Preliminary Report on the Growth of the Rock Bass, *Ambloplites rupestris* (Rafinesque), in Two Lakes of Northern Wisconsin. *Trans. Wis. Acad. Sci., Arts and Letters*, 24: 581-595.

SOUTH LEADS IN THE MILL USE OF COTTON

Fifty years ago, only one-fifth of the cotton spun in the United States was consumed in mills located in the cotton-growing States.

In recent years, however, more than four-fifths of the total mill consumption in the United States has been in the cotton-growing States, according to the Bureau of Agricultural Economics of the United States Department of Agriculture. The cotton-growing States now have more than 70 per cent of the cotton spindles in use in the United States.

Although practically all kinds of cotton goods are produced by southern mills, the proportion of coarse and medium fabrics is larger than in other States. Mills in New England are noted for their output of fine yarns and fabrics.